

Combinatorial Laplacians and Positivity Under Partial Transpose

Roland Hildebrand*

LJK, Université Joseph Fourier, Tour IRMA,
51 rue des Mathématiques, 38400 St. Martin d'Hères, France

Stefano Mancini[†]

Dipartimento di Fisica, Università di Camerino,
Via Madonna delle Carceri, 9, I-62032 Camerino, Italy

Simone Severini[‡]

Department of Mathematics and Department of Computer Science,
University of York, YO10 5DD York, United Kingdom

February 1, 2008

Abstract

Density matrices of graphs are combinatorial laplacians normalized to have trace one (Braunstein *et al. Phys. Rev. A*, **73**:1, 012320 (2006)). If the vertices of a graph are arranged as an array, then its density matrix carries a block structure with respect to which properties such as separability can be considered. We prove that the so-called degree-criterion, which was conjectured to be necessary and sufficient for separability of density matrices of graphs, is equivalent to the PPT-criterion. As such it is not sufficient for testing the separability of density matrices of graphs (we provide an explicit example). Nonetheless, we prove the sufficiency when one of the array dimensions has length

*roland.hildebrand@imag.fr

[†]stefano.mancini@unicam.it

[‡]ss54@york.ac.uk

two (for an alternative proof see Wu, *Phys. Lett. A* **351** (2006), no. 1-2, 18–22).

Finally we derive a rational upper bound on the concurrence of density matrices of graphs and show that this bound is exact for graphs on four vertices.

1 Introduction

Introduced by Braunstein *et al.* [3, 4], density matrices of graphs are simply combinatorial laplacians normalized to have unit trace (the normalization consists of dividing the nonzero entries by twice the cardinality of the edge set). In this way, to any graph G (with labeled vertices) can be associated a specific mixed quantum state (identified with its matrix representation), which is then called the *density matrix of G* . If the vertices of a graph are arranged in a multi-dimensional array, then the density matrix of the graph carries a block structure, which can be associated with a split of the quantum system into subsystems. Each array dimension will then correspond to one subsystem, and the length of the array dimension will equal the number of pure states the subsystem can assume. It is useful to remark that the combinatorial properties of the graph G up to isomorphism do not always characterize its density matrix and therefore do not specify the physical properties of the state. This explains why we need to consider labeled graphs. In other words, we assume that graphs with different adjacency matrices (even if belonging to the same isomorphism class) have different density matrices and then correspond to different quantum states, whose properties can be radically different.

Studying density matrices of graphs with the tool-box provided by quantum mechanics has a twofold role: from the perspective of combinatorics, this interface can be fruitful in uncovering and re-defining graph theoretic properties; from the perspective of quantum mechanics, density matrices of graphs can be seen as “simple” and “highly symmetric” states. Observed under this light, density matrices of graphs provide a restricted testing ground for better understanding techniques and concepts employed in more general settings. Such an approach has particular value, when considering the particular kind of developments in quantum physics and its applications that we are experiencing today.

Indeed, the study of finite-dimensional states is important in quantum in-

formation processing. This is the multidisciplinary area whose goal is to understand and exploit the information encoded in quantum states (see Nielsen and Chuang [20] for a monograph on the subject; see Alber *et al.* [1] for a collection of overviews). The basis of this field consists of interpreting certain quantum physical entities as information carriers and their evolution in time as information processing dynamics. Such a view is giving rise to a number of discoveries and successful real world applications, the most popularized examples being quantum communication and quantum computing. The main ingredient which most likely is responsible for the “quantumness” is the concept of *entanglement*, a property associated to certain quantum states.

While at the early stages of quantum physics entanglement was considered a mystery, nowadays it is recognized as a precious resource, difficult to create and to preserve. Defining entanglement is not an easy task (Bruß [5] compiles an eloquent list of definitions). Speaking about entanglement makes sense only if one considers a system composed of at least two subsystems. The rough idea is that if the two parties (or, equivalently, subsystems) are *entangled* then a complete description of the whole system does not imply a complete description of the parts and *vice versa*. So, two entangled systems present some kind of *correlation* that does not appear to occur in the realm of classical mechanics, where complete information on the system implies a complete description of its individual parts.

From the mathematical point of view, the theory of entanglement is rich and diversified. It has branches in geometry, knots, Lie groups, positive maps, combinatorics, convex optimization, *etc.* The main problems are: *(i)* determine whether a given quantum state is entangled; *(ii)* determine how much entanglement is in a given quantum state; *(iii)* determine the “quality” of entanglement (*e.g.*, the problem of distillability).

As we mentioned above, density matrices of graphs are a restricted set in which these tasks can have a special treatment. Specifically, Braunstein *et al.* [3, 4] and Wu [29] considered the *Quantum Separability Problem (QSP)* for these matrices. The QSP is the computational problem of deciding whether a given quantum state is entangled or not, that is *separable* (Ioannu 2006 [16] is a recent review). The QSP is equivalent to an instance of a combinatorial optimization problem called the Weak Membership Problem and defined in Grötschel *et al.* [10]. In its complete generality the QSP is NP-hard (Gurvits [11]).

There is some evidence that the QSP for density matrices of graphs might

be easier than for general density matrices. A simple necessary condition for separability is that the degrees of the vertices of G are the same as the degrees of the vertices of another graph, G' , obtained from G by mean of a simple operation acting on the edges. The operation is a combinatorial analogue of the linear algebraic partial transposition. In fact, the graph G' will be called here *partially transposed graph*. The condition for separability will be called here *degree-criterion*. Since the partial transposition is centrally involved in the famous *Peres-Horodeckis' criterion* for separability of general states [21, 14] (also called *Positivity under Partial Transpose Criterion*, or, for short, *PPT-criterion*), it is natural to investigate the relation between the degree-criterion and the PPT-criterion when applied to density matrices of graphs.

In this paper, we give an elementary proof that the two criteria are equivalent for density matrices of graphs. We also exhibit a simple example showing that the degree-criterion is not sufficient for testing separability of density matrices of disconnected graphs (that is, graphs with more than one connected component). Additionally, we verify the sufficiency of the degree-criterion when the dimension of one of the parties is two, therefore giving an alternative proof of a result of Wu [29]. There are four sections in the paper. The above observations are exposed in Section 2, after providing the necessary notions and terminology. Section 3 is devoted to point (ii) above. In particular, we focus on the concurrence. This is a quantity associated to every density matrix and it is strictly larger than zero for entangled states (Hill and Wootters [13]). We derive a simple upper bound on the concurrence of density matrices of graphs and show the exactness of this bound for graphs with four vertices. In Section 4 we draw some conclusions.

2 The degree-criterion and the PPT-criterion for density matrices of graphs

The purpose of this section is to shed further light onto the QSP of density matrices of graphs. First, we state formally the QSP and define the PPT-criterion. We then recall the notion of combinatorial laplacian. We define the degree-criterion and we prove its equivalence with the PPT-criterion. We conclude the section by showing that the degree-criterion is necessary and sufficient for testing separability of density matrices of graphs in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^q$

(see also Wu [29]). Our reference on graph theory is Godsil and Royle [9].

2.1 The quantum separability problem

In the axiomatic formulation of quantum mechanics in Hilbert space, the state of a quantum mechanical system, associated to the n -dimensional Hilbert space $\mathcal{H} \cong \mathbb{C}^n$, is identified with an $n \times n$ positive semidefinite, trace-one hermitian matrix, called a *density matrix*. In Dirac notation, a unit vector in a Hilbert space $\mathcal{H} \cong \mathbb{C}^n$ is denoted by $|\psi\rangle$, where ψ is simply a label; given the vectors $|\varphi\rangle, |\psi\rangle \in \mathcal{H}$, the linear functional sending $|\psi\rangle$ to the inner product $\langle\varphi|\psi\rangle$ is denoted by $\langle\varphi|$. (We could easily avoid the Dirac notation here, but we use it to be coherent with the standard literature.) For any unit vector $|\psi\rangle \in \mathcal{H}$, the projector on $|\psi\rangle$ is the hermitian matrix $P[|\psi\rangle] := |\psi\rangle\langle\psi|$, which is called *pure state*. Every density matrix can be written as a weighted sum of pure states (with real nonnegative weights summing up to 1); if the sum has more than one component then the state is said to be *mixed*. By this definition, the decomposition of a mixed density matrix into pure states is not necessarily unique. A matrix of the form $P[|\psi\rangle] \otimes P[|\varphi\rangle]$ is called a *product state*, where the symbol “ $\otimes$ ” denotes the Kronecker or tensor product. Let S_A and S_B be two quantum mechanical systems associated to the p -dimensional and q -dimensional Hilbert spaces $\mathcal{H}_A \cong \mathbb{C}_A^p$ and $\mathcal{H}_B \cong \mathbb{C}_B^q$, respectively. The composite system S_{AB} , which consists of the subsystems S_A and S_B , is associated to the Hilbert space $\mathcal{H}_{AB} \cong \mathbb{C}_A^p \otimes \mathbb{C}_B^q$. The density matrix ρ of S_{AB} is said to be *separable* if

$$\rho = \sum_{i=1}^N p_i P[|\psi_i\rangle_A] \otimes P[|\varphi_i\rangle_B],$$

where $p_i \geq 0$, for every $i = 1, 2, \dots, N$, and $\sum_{i=1}^N p_i = 1$;

the projectors $P[|\psi_i\rangle_A]$ and $P[|\varphi_i\rangle_B]$ are product states acting on $\mathcal{H}_A$ and $\mathcal{H}_B$, respectively. A density matrix ρ is said to be *entangled* if it is not separable. Entangled states cannot be prepared from separable states by mean of operations acting locally on the subsystems. Although the definition given here refers to exactly two parties, entanglement can be defined as well for systems composed of many subsystems.

2.2 The PPT-criterion

The PPT-criterion is based on the notion of partial transpose. This is a common and important notion in the study of entanglement. Let ρ be a density matrix acting on the Hilbert space $\mathcal{H}_{AB} \cong \mathbb{C}_A^p \otimes \mathbb{C}_B^q$. Let

$$\{|u_1\rangle, |u_2\rangle, \dots, |u_p\rangle\} \quad \text{and} \quad \{|w_1\rangle, |w_2\rangle, \dots, |w_q\rangle\}$$

be orthonormal bases of $\mathbb{C}_A^p$ and $\mathbb{C}_B^q$, respectively. Let $\{|v_1\rangle, |v_2\rangle, \dots, |v_n\rangle\}$ be an orthonormal basis of $\mathcal{H}_{AB}$, where $n = pq$. Alternatively, we index these basis vectors with pairs (k, l) . These vectors are taken as follows:

$$|v_{(k-1)q+l}\rangle = |v_{k,l}\rangle = |u_k\rangle |w_l\rangle, \quad k = 1, \dots, p; \quad l = 1, \dots, q.$$

The *partial transpose* of ρ with respect to the system S_B is the $pq \times pq$ matrix, denoted by ρ^{Γ_B} , with the $(i, j; i', j')$ -th entry defined as follows:

$$[\rho^{\Gamma_B}]_{i,j;i',j'} = \langle u_i | \langle w_{j'} | \rho | w_j \rangle | u_{i'} \rangle,$$

where $1 \leq i, i' \leq p$ and $1 \leq j, j' \leq q$. The density matrix of S_{AB} can be written as

$$\rho = \begin{pmatrix} A_{11} & \dots & A_{1p} \\ \vdots & \ddots & \vdots \\ A_{p1} & \dots & A_{pp} \end{pmatrix}, \quad (1)$$

with $q \times q$ matrices A_{ij} acting on the space $\mathbb{C}_B^q$. The partial transpose is then realized by transposing all these matrices:

$$\rho^{\Gamma_B} = \begin{pmatrix} A_{11}^T & \dots & A_{1p}^T \\ \vdots & \ddots & \vdots \\ A_{p1}^T & \dots & A_{pp}^T \end{pmatrix}.$$

If ρ is separable then $\rho^{\Gamma_B} \geq 0$ (see Peres [21]). However, the converse is not necessarily true, since there exist entangled states with positive partial transpose (the so-called *bound entangled states*). The failure of the PPT-criterion is then the failure of an operational characterization of entangled states, which is computationally simple to verify. The PPT-criterion is necessary and sufficient for separability of density matrices acting on $\mathbb{C}_A^2 \otimes \mathbb{C}_B^2$ or $\mathbb{C}_A^2 \otimes \mathbb{C}_B^3$ (Horodecki *et al.* [14]); it is also necessary and sufficient for certain infinite-dimensional states (Simon [25]; Duan *et al.* [7]; Mancini and

Severini [26] is a brief review). It is important to mention that only one other (operational) criterion is known for detecting entanglement: the *realignment criterion* (Rudolph [23]; Chen [6]). It can detect bound entanglement, but for some states it is weaker than the PPT-criterion. Unfortunately, it can be checked numerically that the two criteria together do not solve the QSP for all states (see Horodecki and Lewenstein [15]). Generally, the operational characterization of entanglement is an open problem.

2.3 Combinatorial laplacians

In this subsection we recall the notion of combinatorial laplacian. A *graph* $G = (V, E)$ is a pair defined in the following way: $V(G)$ is a non-empty and finite set whose elements are called *vertices*; $E(G)$ is a non-empty set of unordered pairs of vertices, which are called *edges*. A *loop* is an edge of the form $\{v_i, v_i\}$, for some vertex v_i . We assume that $E(G)$ does not contain loops. A graph G is said to be *on n vertices* if the number of elements in $V(G)$ is n . The *adjacency matrix* of a graph on n vertices G is an $n \times n$ matrix, denoted by $M(G)$, having rows and columns labeled by the vertices of G , and ij -th entry defined as follows¹:

$$[M(G)]_{i,j} := \begin{cases} 1, & \text{if } \{v_i, v_j\} \in E(G); \\ 0, & \text{if } \{v_i, v_j\} \notin E(G). \end{cases}$$

Two vertices v_i and v_j are said to be *adjacent* if $\{v_i, v_j\} \in E(G)$. The *degree* of a vertex $v_i \in V(G)$, denoted by $d_G(v_i)$, is the number of edges adjacent to v_i . The *degree-sum* of G is defined and denoted by

$$d_G = \sum_{i=1}^n d_G(v_i).$$

The *degree matrix* of G is an $n \times n$ matrix, denoted by $\Delta(G)$, having ij -th entry defined as follows:

$$[\Delta(G)]_{i,j} := \begin{cases} d_G(v_i), & \text{if } i = j; \\ 0, & \text{if } i \neq j. \end{cases}$$

The *combinatorial laplacian matrix* of a graph G (for short, *laplacian*) is the matrix

$$L(G) := \Delta(G) - M(G).$$

¹We are considering here only ‘simple’ graphs.

According to our definition of graph, $L(G) \neq 0$. Moreover, the laplacian of a graph G , scaled by the degree-sum of G , has trace one and is semidefinite positive. As such it has the characteristic features of a quantum mechanical density matrix, hence it would provide a link to quantum states. On the basis of this observation, we fix the following definition: the *density matrix of a graph G* is the matrix

$$\rho_G := \frac{1}{d_G} L(G).$$

Let $\mathcal{G}_n$ be the set of density matrices of graphs on n vertices. The set $\mathcal{G}_n$ is a subset of the set of all density matrices acting on the n -dimensional Hilbert space $\mathcal{H}_{AB} \cong \mathbb{C}_A^p \otimes \mathbb{C}_B^q$, where $n = pq$. The number of elements in $\mathcal{G}_n$ equals the number of graphs on n vertices, a number that grows superexponentially in n . There are many applications of laplacians. In particular, their eigensystems are a rich source of information about graphs (Mohar [19]).

It is important to remark that graphs with different adjacency matrices have different density matrices, even graphs belonging to the same isomorphism class (e.g. those obtained from each other by permutation of the vertices labels). In fact, given a graph G with density matrix ρ_G , if there exist a permutation matrix P such that $P^T M(G) P = M(G')$, then $G \cong G'$. As a consequence G' has density matrix $P^T \rho_G P$.

Finally, given the density matrix ρ_G of a graph, in order to have a correspondence to a quantum state (density operator), we have to specify the basis in the Hilbert space with respect to which the quantum state (density operator) has ρ_G as matrix representation. This can be done by associating vertices labels to orthonormal vectors.

2.4 The degree-criterion

Let G be a graph on $n = pq$ vertices $v_1, v_2, \dots, v_n$. These vertices are represented here as ordered pairs in the following way:

$$v_{(k-1)p+l} = (u_k, w_l) \equiv u_k w_l, \quad k = 1, \dots, p; \quad l = 1, \dots, q.$$

By respecting this labelling, we associate G to the orthonormal basis

$$\{|v_i\rangle : i = 1, 2, \dots, n\} = \{|u_j\rangle \otimes |w_k\rangle : j = 1, 2, \dots, p; k = 1, 2, \dots, q\},$$

of the Hilbert space $\mathcal{H}_{AB} \cong \mathbb{C}_A^p \otimes \mathbb{C}_B^q$, where

$$\{|u_j\rangle : j = 1, 2, \dots, p\} \quad \text{and} \quad \{|w_k\rangle : k = 1, 2, \dots, q\}.$$

are orthonormal bases of the Hilbert spaces $\mathcal{H}_A \cong \mathbb{C}^p$ and $\mathcal{H}_B \cong \mathbb{C}^q$, respectively. The *partial transpose of a graph* $G = (V, E)$ (with respect to $\mathcal{H}_B$), denoted by $G^{\Gamma_B} = (V, E')$, is the graph such that

$$\{u_i w_j, u_k w_l\} \in E' \quad \text{if and only if} \quad \{u_i w_l, u_k w_j\} \in E.$$

If $\Delta(G) = \Delta(G^{\Gamma_B})$ we say that G satisfies the *degree-criterion*. The following conjecture was proposed in Braunstein *et al.* [4]: a density matrix ρ_G of a graph on $n = pq$ vertices is separable in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ if and only if $\Delta(G) = \Delta(G^{\Gamma_B})$. A proof of this conjecture would give a simple method for testing the separability of density matrices of graphs, as we would only need to check whether the $n \times n$ diagonal matrices $\Delta(G)$ and $\Delta(G^{\Gamma_B})$ are equal. There are counterexamples to this conjecture, when the graph has isolated vertices (that is, vertices not belonging to any edge). This is the case for the graph G defined on a 3×3 array, with laplacian

$$L(G) = \begin{pmatrix} I_4 & 0 & -I_4 \\ 0 & \vdots & 0 \\ -I_4 & 0 & I_4 \end{pmatrix},$$

where I_d is the d -dimensional identity matrix. Indeed, G satisfies the degree-criterion, but ρ_G is entangled. For connected graphs, we don't have any counterexample yet. Next, we list known partial results about separability:

- Let ρ_G be the density matrix of a graph on $n = pq$ vertices. If ρ_G is separable in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ then $\Delta(G) = \Delta(G^{\Gamma_B})$ (Braunstein *et al.* [4]).
- Let G be a nearest point graph on $n = pq$ vertices. Then the density matrix ρ_G is separable in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ if and only if $\Delta(G) = \Delta(G^{\Gamma_B})$ (Braunstein *et al.* [4]). (It may be worth recalling the definition of nearest point graph. Consider a rectangular lattice with pq points arranged in p rows and q columns, such that the distance between two neighboring points on the same row or in the same column is 1. A *nearest point graph* is a graph whose vertices are identified with the points of the lattice and the edges have length 1 or $\sqrt{2}$.)
- Let G and H be two graphs on $n = pq$ vertices. If ρ_G is separable in $\mathbb{C}^p \otimes \mathbb{C}^q$ and $G \cong H$ (that is, G and H are isomorphic) then ρ_H is not necessarily separable in $\mathbb{C}^p \otimes \mathbb{C}^q$. However, there are exceptions, as observed by the following point (Braustein *et al.* [3]).

- Let K_n be the *complete graph* on n vertices. Recall that the complete graph is the graph with an edge between any pair of vertices. One can show that, for any $n = pq$, the density matrix ρ_{K_n} is separable in $\mathbb{C}^p \otimes \mathbb{C}^q$. Notice that for a graph H such that $M(H) = M(G) \otimes M(G')$ for some graphs G and G' , the density matrix ρ_H is separable. Of course, if a density matrix ρ_G is separable it does not necessarily mean that $M(G)$ is a tensor product. The *star graph* on n vertices $v_1, v_2, \dots, v_n$, denoted by $K_{1,n-1}$, is the graph whose set of edges is $\{\{v_1, v_i\} : i = 2, 3, \dots, n\}$. The density matrix $\rho_{K_{1,n-1}}$ is entangled for $n = pq \geq 4$. So, the separability properties of complete graphs and star graphs do not depend on the labelling. It is an open problem to determine if these graphs are the only ones with this property (Braunstein *et al.* [3]).
- Let ρ_G be the density matrix of a graph on $n = 2q$ vertices. Then $\rho_G^{\Gamma_B} \geq 0$ if and only if ρ_G is separable in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^q$. Equivalently, the PPT-criterion is necessary and sufficient to test separability in the described case (Wu [29]).
- Wu [29] considered generalized laplacians. Let S be the set of density matrices with nonnegative row sums and nonpositive off-diagonal entries. If a density matrix $\rho \in S$ of dimension $n = pq$ is such that the matrices A_{ij} (as in Eq. 1) are line sum symmetric, then ρ is separable in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$. A matrix is *line sum symmetric* if the i -th column sum is equal to the i -th row sum for each i . As a corollary, Wu [29] proved that, if a density matrix $\rho \in S$ of dimension $n = pq$ and with zero row sums is such that $[\rho]_{i,j;i',j'} \neq 0$ implies that $|i - i'| \leq 1$ then ρ is separable in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ if and only if ρ^{Γ_B} has zero row sums (Corollary 3). This result generalizes separability of nearest point graphs. In fact, for a nearest point graph also the condition $|j - j'| \leq 1$ is required. It is relevant to notice that ρ^{Γ_B} has zero row sum if and only if the degree-criterion is satisfied.

2.5 Equivalence of degree and PPT-criterion

Here we prove that for laplacians the PPT-criterion is equivalent to the degree criterion.

Observation 1. Let ρ be a matrix acting on $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ and satisfying the PPT-criterion. Let $x \otimes y$ be a separable vector in $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$. Then the

condition $\rho(x \otimes y) = 0$ implies the condition $\rho^{\Gamma_B}(x \otimes \bar{y}) = 0$. In fact, we have $(x \otimes y)^* \rho(x \otimes y) = (x \otimes \bar{y})^* \rho^{\Gamma_B}(x \otimes \bar{y}) = 0$, and by the positivity of ρ^{Γ_B} it follows that $\rho^{\Gamma_B}(x \otimes \bar{y}) = 0$.

Here the star denotes adjoint and the overbar denotes complex conjugation. A simple proof of Theorem 2 by Braunstein *et al.* [4] can be derived from this result, with x and y being equal to the all-ones vector.

Observation 2. As a consequence of Observation 1, if ρ is a separable density matrix and

$$\rho = \sum_{k=1}^N (x_k \otimes y_k)(x_k \otimes y_k)^*$$

is a separable decomposition of ρ then, for any $k = 1, 2, \dots, N$, we have the following conditions:

- $(x_k \otimes y_k) \in \text{range}(\rho)$;
- $(x_k \otimes \bar{y}_k) \in \text{range}(\rho^{\Gamma_B})$.

Theorem 1. Let ρ_G be the density matrix of a graph G . Then ρ_G satisfies the PPT-criterion if and only if it satisfies the degree-criterion.

Proof. We have $\rho_G(e \otimes e) = 0$, because ρ_G is the laplacian of G scaled by some coefficient, where e is the all-ones vector of the required dimension. Suppose that the degree-criterion is satisfied. Then $\rho_G^{\Gamma_B} = \rho_{G^{\Gamma_B}}$. Hence $\rho_G^{\Gamma_B}$ is positive. It follows that ρ_G satisfies the PPT-criterion. Suppose that the PPT-criterion is satisfied. Then, by Observation 1, we have that $\rho_G^{\Gamma_B}(e \otimes e) = 0$. This is exactly the degree-criterion on ρ_G . ■

2.6 Separability in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^q$

Here we prove that the degree-criterion is necessary and sufficient to test separability in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^q$ of density matrices of graphs, therefore giving an alternative proof to a result of Wu [29].

Theorem 2. Let G be a graph on $n = 2q$ vertices. Then ρ_G is separable in $\mathbb{C}_A^2 \otimes \mathbb{C}_B^q$ if and only if the degree-criterion is satisfied.

Proof. The implication “ $\Rightarrow$ ” is easily verified. We prove the implication “ $\Leftarrow$ ”. If G satisfies the degree criterion then we can write

$$\rho_G = L_1 + L_2 + L_3,$$

where

$$L_1 := \begin{pmatrix} X_1 & 0 \\ 0 & 0 \end{pmatrix}, L_2 := \begin{pmatrix} 0 & 0 \\ 0 & X_2 \end{pmatrix} \text{ and } L_3 := \begin{pmatrix} X_3 & X_4 \\ X_4^T & X_3 \end{pmatrix}$$

and $X_1, \dots, X_4$ are appropriate matrices. Now, L_1 and L_2 are trivially separable. The matrix L_3 is separable because it is a PSD block-Töplitz matrix. Hence, ρ_G is separable. ■

3 Concurrence

In this section, we focus on the concurrence of density matrices of graphs. The notion of concurrence was introduced by Hill and Wootters [13]. The concurrence of a density matrix acting on $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$ is a quantity which is strictly larger than zero if the state is entangled and zero if it is separable. Here is the definition. Let $|\psi\rangle_{AB} \in \mathbb{C}_A^p \otimes \mathbb{C}_B^q$. The *concurrence* of $|\psi\rangle_{AB}$ is denoted and defined as follows:

$$\mathcal{C}(\psi) = \sqrt{2(1 - \text{tr}(\rho_A^2))},$$

where

$$\rho_A = \text{tr}_B(|\psi\rangle_{AB}\langle\psi|).$$

Let ρ_{AB} be a density matrix acting on $\mathbb{C}_A^p \otimes \mathbb{C}_B^q$. The concurrence of ρ_{AB} is denoted and defined as

$$\mathcal{C}(\rho_{AB}) = \inf \left\{ \sum_i \omega_i \mathcal{C}(\psi_i) : \rho_{AB} = \sum_i \omega_i |\psi_i\rangle_{AB}\langle\psi_i|, 0 \leq \omega_i \leq 1, \sum_i \omega_i = 1 \right\}.$$

Let now $p = q = 2$ and

$$\sigma_y = -i|1\rangle\langle 2| + i|2\rangle\langle 1|,$$

where $|1\rangle$ and $|2\rangle$ are the eigenvectors of the matrix

$$\sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

corresponding to the eigenvalues 1 and -1 , respectively. An analytical formula for $\mathcal{C}(\rho_{AB})$, is given by

$$\mathcal{C}(\rho_{AB}) = \max\{0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4\},$$

where $\lambda_1, \lambda_2, \lambda_3$ and λ_4 are the square roots of the eigenvalues of $\rho_{AB}\tilde{\rho}_{AB}$ arranged in decreasing order and

$$\tilde{\rho}_{AB} := (\sigma_y \otimes \sigma_y) \rho_{AB}^T (\sigma_y \otimes \sigma_y).$$

The importance of the concurrence stems from its relation with the so-called *entanglement of formation*, the most widely accepted measure of entanglement (Bennett *et al.* [2]; see also Plenio and Virmani [22]). For a pure state (that is a state of the form $P[|\psi\rangle]$) of a system S_{AB} , a good measure of entanglement is the entropy of the density matrix associated with one of the two subsystems. Choosing the system S_A , this can be written as

$$E(\psi) := -\text{tr}(\rho_A \log_2 \rho_A),$$

where

$$\rho_A = \text{tr}_B(P[|\psi\rangle]).$$

For a mixed state ρ , the entanglement of formation is defined by

$$E_f(\rho) := \min \sum_i p_i E(\psi_i),$$

where the minimum is taken over all pure states decompositions of the density matrix ρ . It is evident that computing E_f is in general not an easy task. Explicit formulas are only known for very specific classes of states. An example consists of Werner states (Vollbrecht and Werner [28]). The role of concurrence is explained by the following result (Wootters [30]). Let ρ be a mixed density matrix of dimension 4. Then


$$E_f(\rho) = H\left(\frac{1}{2} + \frac{1}{2}\sqrt{1 - \mathcal{C}(\rho)^2}\right),$$

where

$$H(x) = -x \ln x - (1 - x) \ln(1 - x),$$

is the standard information-theoretic entropy. Remarkably, $E_f(\rho)$ increases monotonically as a function of $\mathcal{C}(\rho)$.

For density matrices of graphs of dimension 4 the situation can be described as follows. There are twelve nonisomorphic graphs on 4 vertices. Seven of these graphs can have entangled density matrices. The tables below present these graphs and their respective concurrence:



Notice that in all cases the value of the concurrence is 1 over the number of edges. Moreover, one easily sees that the optimal decomposition of the density matrices of these graphs just corresponds to the decomposition of the combinatorial laplacians as sums of laplacians of 1-edge graphs.

This motivates the following definitions.

Definition 1 Let $G = (V, E)$ be a graph, with its $n = pq$ vertices arranged in a $p \times q$ array. We call an edge $e \in E$ separable if the density matrix of the 1-edge graph $G_e = (V, \{e\})$ is separable. We call $e \in E$ matched if $e \in E'$ and unmatched otherwise.

Recall that E' is the set of edges of the partially transposed graph G' . Thus an edge is matched if and only if it is part of a criss-cross² or it is separable. Since graphs consisting of a criss-cross give rise to separable density matrices, we have the following results.

Observation 3. Let $G = (V, E)$ be as above. Let further $E_1 \subset E$ be the subset of all matched edges. Then the graph $G_1 = (V, E_1)$ has a separable density matrix.

² A *criss-cross* is a set $\{\{(k, i), (l, j)\}, \{(k, j), (l, i)\}\}$ of two edges belonging to a set of (vertex-disjoint) entangled edges on $n = pq$ vertices (see also [3]).

Observation 4. Let $G = (V, \{e\})$ be a 1-edge graph, with its $n = pq$ vertices arranged in a $p \times q$ array, and let ρ_e be its density matrix. Then the concurrence of ρ_e is given by 0 if e is a separable edge and by 1 if e is not separable, and hence unmatched.

Corollary 2 *Let $G = (V, E)$ be as above. Let n_1 be the number of matched edges of G and n_2 be the number of unmatched edges. Then the concurrence of the density matrix ρ of G is bounded from above by*

$$\mathcal{C}(\rho) \leq \frac{n_2}{n_1 + n_2}.$$

In particular, for any graph G with density matrix ρ we have $\mathcal{C}(\rho) \leq 1$.

Proof. Assume the above notations. Let ρ_1, ρ_2 be the density matrices of the graphs (V, E_1) and (V, E_2) , respectively, where E_1 is the set of matched edges and E_2 the set of unmatched edges. Then the density matrix ρ of G is given by the convex combination $\rho = \frac{n_1}{n_1 + n_2} \rho_1 + \frac{n_2}{n_1 + n_2} \rho_2$ and the density matrix ρ_2 is given by $\rho_2 = \frac{1}{n_2} \sum_{e \in E_2} \rho_e$, where ρ_e is the density matrix of the 1-edge graph $(V, \{e\})$. By convexity of the concurrence and by Observations 3 and 4 we obtain

$$\mathcal{C}(\rho) \leq \frac{n_1}{n_1 + n_2} \mathcal{C}(\rho_1) + \sum_{e \in E_2} \frac{1}{n_1 + n_2} \mathcal{C}(\rho_e) = \frac{n_2}{n_1 + n_2}.$$

■

For all of the above graphs on four vertices either the density matrix is separable or we have $n_2 = 1$, in which case 1 over the number of edges is an upper bound for the concurrence. As can be seen from the table, the bound is actually achieved.

The concurrences of graph density matrices which have rank 2 are listed in [12]. Examples IVb, IVc and IX in [12] or the tally-mark³ show that in general the upper bound is not exact, even for graphs on 2×3 arrays.

³ A *tally-mark* is a set

$$\{(k, i_1), (l, i_2)\}, \{(k, i_2), (l, i_3)\}, \dots, \{(k, i_{s+1}), (l, i_{s+2})\}, \{(k, i_{s+2}), (l, i_1)\}$$

of $s + 2$ edges, where $k < l$, $s \geq 0$ and $i_1 < i_2 < \dots < i_{s+2}$, belonging to a set of (vertex-disjoint) entangled edges on $n = pq$ vertices (see also [3]). Note that a criss-cross is a tally-mark with two edges.

Definition 3 We call a graph $G = (V, E)$ maximally entangled if the concurrence of its density matrix ρ is given by $\mathcal{C}(\rho) = 1$.

All edges of a maximally entangled graph must hence be unmatched.

Some natural questions arise:

- Are there nonisomorphic graphs with the same concurrence?
- How can graphs with rational concurrence be characterized?
- Is the concurrence of ρ_G related to specific combinatorial properties of G ?
- Can the set of edges of a graph $G = (V, E)$ with density matrix ρ always be partitioned in two subsets E_1, E_2 such that the density matrix of (V, E_1) is separable, (V, E_2) is maximally entangled and $\mathcal{C}(\rho) = \frac{\#E_2}{\#E}$?
- How can maximally entangled graphs be characterized?
- Does there exist a class of density matrices of graphs for which testing separability is a difficult problem? Maybe, the existence of such a class would provide a transparent proof that detecting entanglement is hard.

Unfortunately, explicit formulae for computing the concurrence of density matrices are known so far only for dimensions $n \leq 4$ (see, Rungta *et al.* [24], Li and Zhu [17], and Mintert *et al.* [18]) and for density matrices of rank 2 [12]. This is an obstacle when thinking about the questions above. Nevertheless, one can still hope to find an *ad-hoc* formula for $\mathcal{C}(\rho)$, when ρ is the density matrix of a graph. In fact, it may well be that the optimal decompositions of ρ_G in pure states are very special. Finding such a formula would be interesting in view of potential generalizations.

Putting the concurrence on a side, one may ask if there is some entanglement measure specifically tailored for ρ_G . Considering the apparent success of the degree-criterion, a naive measure would be the normalized Euclidean norm $EN(\rho_G) := \|\Delta(G) - \Delta(G^{\Gamma_B})\|$. The *logarithmic negativity* is a well-known entanglement measure and it is defined by $LN(\rho_G) := \log_2(1 + 2\mathcal{N}(\rho_G))$, where $\mathcal{N}(\rho_G)$ is the sum of the magnitudes of all negative eigenvalues of $\rho_G^{\Gamma_B}$ (see Vidal and Werner [27]). There are examples of graphs G and H for which $EN(\rho_G) = EN(\rho_H)$ but $LN(\rho_G) \neq LN(\rho_H)$ (Ghosh [8]).

4 Conclusion

We have proven that the degree-criterion is equivalent to the PPT-criterion. It is thus in general not sufficient for separability of density matrices of graphs. As a matter of fact, we have provided a counterexample within graphs having isolated vertices. Nevertheless, we have been able to prove the sufficiency of the degree-criterion when one of the subsystems has dimension two. In particular, as a corollary of Theorem 2, one can easily obtain the separability of criss-crosses and tally-marks.

We have also considered the concurrence as a possible entanglement measure of density matrices of graphs. There could be more suitable entanglement measures for such kind of states, especially because no explicit formula for concurrence is known when $n > 4$ and the rank of the density matrix is exceeding 2. Further studies are in order on the subject of density matrices of graphs. However we believe that such states provide a restricted testing ground for better understanding techniques and concepts employed in more general settings.

References

- [1] G. Alber *et al.* (Eds.), *Quantum Information: An Introduction to Basic Theoretical Concepts and Experiments*, Series: Springer Tracts in Modern Physics, Vol. 173, 2001.
- [2] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Mixed-state entanglement and quantum error correction, *Phys. Rev. A* (3) **54** (1996), no. 5, 3824–3851, quant-ph/9604024.
- [3] S. L. Braunstein, S. Ghosh and S. Severini, The laplacian of a graph as a density matrix: a basic combinatorial approach to separability of mixed states, to appear in *Ann. Comb.*, quant-ph/0406165.
- [4] S. L. Braunstein, S. Ghosh, T. Mansour, S. Severini, and R. C. Wilson, Some families of density matrices for which separability is easily tested, *Phys. Rev. A*, **73**:1, 012320 (2006), quant-ph/0508020.
- [5] D. Bruß, Characterizing entanglement. Quantum information theory. *J. Math. Phys.* **43** (2002), no. 9, 4237–4251. quant-ph/0110078.

- [6] K. Chen and L.-A. Wu, A matrix realignment method for recognizing entanglement, *Quantum Inf. Comput.* **3** (2003), no. 3, 193–202, quant-ph/0205017.
- [7] L. M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, Inseparability criterion for continuous variable systems, *Phys. Rev. Lett.* **84**, 2722 (2000), quant-ph/9908056.
- [8] S. Ghosh, Personal communication, June 2006.
- [9] C. Godsil and G. Royle, Algebraic graph theory. Graduate Texts in Mathematics, 207. *Springer-Verlag, New York*, 2001.
- [10] M. Grötschel, L. Lovász, and A. Schrijver, *Geometric Algorithms and Combinatorial Optimization* (Springer-Verlag, Berlin, 1988).
- [11] L. Gurvits, Classical deterministic complexity of Edmonds’ Problem and quantum entanglement, *Proceeding of the thirty-fifth ACM Symposium on Theory of Computing (ACM Press, New York)*, 2003, 10-19, quant-ph/0303055.
- [12] R. Hildebrand, Concurrence of Lorentz-positive maps, quant-ph/0612064.
- [13] S. Hill and W. K. Wootters, Entanglement of a Pair of Quantum Bits, *Phys. Rev. Lett.* **78**, 5022 (1997).
- [14] M. Horodecki, P. Horodecki, and R. Horodecki, Separability of mixed states: necessary and sufficient conditions, *Phys. Lett. A* **223** (1996), no. 1-2, 1–8, quant-ph/9605038.
- [15] P. Horodecki and M. Lewenstein, Bound entanglement and continuous variables, *Phys. Rev. Lett.* **85** (2000), no. 13, 2657–2660, quant-ph/0001035.
- [16] L. M. Ioannou, Deterministic Computational Complexity of the Quantum Separability Problem, quant-ph/0603199.
- [17] Y.-Q. Li, G.-Q. Zhu, Concurrence Vectors for Entanglement of High-dimensional Systems, quant-ph/0308139.

- [18] F. Mintert, M. Kuś, and A. Buchleitner, Concurrence of mixed multipartite quantum states, *Phys. Rev. Lett.* **95** (2005), no. 26, 260502, quant-ph/0411127.
- [19] B. Mohar, *The Laplacian spectrum of graphs*. Graph theory, combinatorics, and applications. Vol. 2 (Kalamazoo, MI, 1988), 871—898, Wiley-Intersci. Publ., Wiley, New York, 1991.
- [20] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [21] A. Peres, Separability criterion for density matrices, *Phys. Rev. Lett.* **77** (1996), no. 8, 1413–1415, quant-ph/9604005.
- [22] M. B. Plenio and S. Virmani, An introduction to entanglement measures, quant-ph/0504163.
- [23] O. Rudolph, Further results on the cross norm criterion for separability, *Quantum Inf. Process.* **4** (2005), no. 3, 219–239, quant-ph/0202121.
- [24] P. Rungta, V. Bužek, Carlton M. Caves, M. Hillery, and G. J. Milburn, Universal state inversion and concurrence in arbitrary dimensions, *Phys. Rev. A* (3) **64** (2001), no. 4, 042315, quant-ph/0102040.
- [25] R. Simon, Peres-Horodecki Separability Criterion for Continuous Variable Systems, *Phys. Rev. Lett.* **84**, 2726–2729 (2000), quant-ph/9909044.
- [26] S. Mancini and S. Severini, The Quantum Separability Problem for Gaussian States, *LMCS06 (Logic, models and computer science)*, Camerino, Italy, to appear in *ENTCS*, cs.CC/0603047.
- [27] G. Vidal and R. Werner, Computable measure of entanglement, *Phys. Rev. A* **65**, 032314 (2002).
- [28] K. G. H. Vollbrecht and R. F. Werner, Entanglement measures under symmetry, *Phys. Rev. A* **64**, 062307 (2001), quant-ph/0010095.
- [29] C. W. Wu, Conditions for separability in generalized Laplacian matrices and diagonally dominant matrices as density matrices, *Phys. Lett. A* **351** (2006), no. 1-2, 18–22, quant-ph/0508163.

- [30] W. K. Wootters, Entanglement of Formation of an Arbitrary State of Two Qubits, *Phys. Rev. Lett.* **80**, 2245 (1998).